

University of Groningen

Two homologous oligopeptide binding protein genes (oppA) in *Lactococcus lactis* MG1363

Sanz, Yolanda; Lanfermeijer, Frank C.; Hellendoorn, Michiel; Kok, Jan; Konings, Wil N.; Poolman, Bert

Published in:
International Journal of Food Microbiology

DOI:
[10.1016/j.ijfoodmicro.2004.04.003](https://doi.org/10.1016/j.ijfoodmicro.2004.04.003)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2004

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Sanz, Y., Lanfermeijer, F. C., Hellendoorn, M., Kok, J., Konings, W. N., & Poolman, B. (2004). Two homologous oligopeptide binding protein genes (oppA) in *Lactococcus lactis* MG1363. *International Journal of Food Microbiology*, 97(1), 9 - 15. <https://doi.org/10.1016/j.ijfoodmicro.2004.04.003>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Two homologous oligopeptide binding protein genes (*oppA*) in *Lactococcus lactis* MG1363

Yolanda Sanz^{a,*}, Frank C. Lanfermeijer^a, Michiel Hellendoorn^b, Jan Kok^b,
Wil N. Konings^a, Bert Poolman^{a,1}

^aDepartment of Microbiology, Groningen Biomolecular Sciences and Biotechnology Institute, University of Groningen,
Kerklaan 30, NL 9751 NN, Haren, The Netherlands

^bDepartment of Genetics, Groningen Biomolecular Sciences and Biotechnology Institute, University of Groningen, Kerklaan 30,
NL 9751 NN, Haren, The Netherlands

Received 19 June 2003; received in revised form 30 September 2003; accepted 4 April 2004

Abstract

In previous studies, it has been shown that inactivation of *opp* or even *oppA* abolishes the capacity of *Lactococcus lactis* to utilize oligopeptides. We now show that the *opp* operon has been duplicated in *L. lactis* MG1363. The nucleotide sequence of the *oppA* and *oppC* homologues (*appA* and *appC*) and most of the *oppB* homologue (*appB*) indicate that the corresponding protein sequences are 83%, 92% and 91% identical, respectively. Inactivation of *appA*, via homologous recombination, as well as complementation studies were carried out to determine the possible function of *appA* in peptide utilization. As anticipated from studies with an *oppA* knock-out, peptide utilization was not impaired in an *appA* disruption mutant. Importantly, AppA expressed from a plasmid could restore the ability of *oppA* deletion mutants to utilize Leu-enkephalin, albeit with a lower efficiency than OppA. The differences in the ability to utilize this pentapeptide were not due to differences in expression levels but most likely reflect a different catalytic efficiency in oligopeptide utilization when AppA is used as ligand receptor.

© 2004 Elsevier B.V. All rights reserved.

Keywords: *Lactococcus lactis*; Oligopeptide transport system; Gene duplication; Gene complementation

1. Introduction

The proteolytic pathway for the utilization of milk proteins as exogenous nitrogen source is well-de-

scribed in *Lactococcus lactis*. The system is composed of an extracellular proteinase, at least three peptide transport systems, one specific for oligopeptides (Opp) and two others for di- and tripeptides (DtpT and Dpp), and a set of intracellular peptidases (Kunji et al., 1996b; Sanz et al., 2001). The extracellular proteinase degrades caseins predominantly into oligopeptides of 5–26 residues, which are exclusively substrates of Opp. The intracellular peptidases contribute to the complete degradation of oligopeptides into free amino acids. Thus, Opp is considered essential for nutrition, while the di-tripeptide transport

* Corresponding author. Present address: Instituto de Agroquímica y Tecnología de Alimentos (C.S.I.C.). Apartado 73. 46100 Burjasot, Valencia Spain. Tel.: +34-96-3900022; fax: +34-96-3636301.

E-mail address: yolsanz@iata.csic.es (Y. Sanz).

¹ Present address: Department of Biochemistry, Groningen Biomolecular Sciences and Biotechnology Institute, University of Groningen, Nijenborgh 4, 9747 AG Groningen, The Netherlands.

systems seem to have a role in regulation (Marugg et al., 1995; Kunji et al., 1996a; Meijer et al., 1996; Sanz et al., 2001). The Opp system belongs to the ABC-transporter superfamily and consists of a peptide binding protein (OppA), two integral membrane proteins (OppB and OppC) and two ATP-binding proteins (OppD and OppF; Tynkkynen et al., 1993). In general, the solute binding proteins provide the primary interaction site for the ligand and largely define the specificity of the transport system (Sleigh et al., 1997). Some binding-protein dependent ABC-transporters, however, function with more than one ligand binding protein, each with a different specificity (Ehrmann et al., 1998). In Gram-positive bacteria, this is exemplified by the three-homologous binding proteins working together in the Ami system of *Streptococcus pneumoniae* (Alloing et al., 1994). In *L. lactis*, the transport of oligopeptides has been exclusively attributed to the Opp system and the binding protein OppA is considered essential for its function.

Here, we report on the identification and characterization of the genes specifying a putative second oligopeptide transport system in *L. lactis* MG1363. Complete sequences of the *oppA* and *oppC* counterparts (*appA* and *appC*) and part of the *oppB* counterpart (*appB*) are presented. The possible functionality of *appA* in peptide utilization has been determined in a gene knockout and by complementation studies.

2. Materials and methods

2.1. Strains and culture conditions

Escherichia coli strains were grown, at 37 °C, in Luria-Bertani broth (Sambrook et al., 1989). Ampicillin or erythromycin was added to 100 µg ml⁻¹, when appropriate. *L. lactis* strains were routinely grown, at 30 °C, in M17 broth (Difco, East Molesey, UK) supplemented with 0.5% (w/v) glucose. The ability of the *L. lactis* strains to utilize peptides was determined in CDM (Poolman and Konings, 1988) containing all amino acids except an essential one that was supplied in the form of a di-, tri- or oligopeptide, as described elsewhere (Kunji et al., 1993). Erythromycin was added to 5 µg ml⁻¹, when appropriate.

2.2. General DNA manipulation and sequence analysis

Standard molecular techniques were essentially performed as described by Sambrook et al. (1989). Chromosomal and plasmid DNA were isolated by the methods of Birnboim and Doly (1979) and Leenhouts et al. (1990). Southern analysis was carried out by the non-radioactive DNA Labeling and Detection kit (Boehringer, Almere, The Netherlands). *E. coli* and *L. lactis* were transformed by electroporation as described by Zabarovsky and Winberg (1990) and Holo and Nes (1989), respectively. Sequence analyses were carried out by the dideoxy-chain termination method (Sanger et al., 1977). Sequence similarity searches were done with the FASTA algorithm of Pearson and Lipman (1988). Sequence alignments were performed with DNAMAN program (Version 4.03).

2.3. Plasmid construction

The *appA* gene was inactivated in the double peptide transport mutant *L. lactis* AG500 (Δopp , $\Delta dtpT$, Hagting et al., 1994; Kunji et al., 1996b) by plasmid integration via homologous recombination (Leenhouts et al., 1989). A 1.25-kb fragment of the *oppA* homologue (*appA*) was initially detected in a genome library of *L. lactis* MG1363 in *E. coli* NM522 (Buist et al., 1995). The 1.25-kb *appA* fragment in p15D4 (Ap^r, pUC19 derivative) was inserted at the *Hind*III/*Asp*718 sites of the integrative vector pORI28 (Em^r) (Leenhouts and Venema, 1993). *L. lactis* AG500 was transformed with the new construct, p15D14dis, to achieve the inactivation of *appA*, which was verified by Southern hybridization. The expression vector of *appA* was constructed as follows. The gene *appA* was amplified from chromosomal DNA of *L. lactis* AG500 and the PCR product was inserted at *Nco*I/*Bam*HI sites of pAMP31 (Picón et al., 2000), thereby replacing *oppA* for *appA*. In this vector, *appA* is under the control of the constitutive lactococcal promoter *P32* (Van der Vossen et al., 1987) and fused to a 3'-end sequence specifying a six-histidine tag. The resulting vector, pYSAprime, was transformed into the *appA* deletion mutant *L. lactis* AMP2 (Picón et al., 2000).

2.4. Western analysis

The expression of *oppA* and *appA* from the corresponding plasmids in whole-cell lysates of *L. lactis* AMP2 was quantified by Western analysis using monoclonal antibodies (Dianova, Apeldoorn, The Netherlands) raised against the 6-histidine tag (Sanz et al., 2000).

2.5. Nucleotide sequence accession number

The sequence data have been submitted to the GenBank database under accession number AF245305.

3. Results and discussion

3.1. Identification of *opp* homologues and sequence analysis

In the chromosome of *L. lactis* MG1363 the existence of a DNA fragment homologous to *oppA* was revealed by Southern analysis (data not shown). Subsequently, a gene library of *L. lactis* MG1363 in *E. coli* NM522 (Buist et al., 1995) was screened for hybridization with a probe consisting of a 750-bp *oppA* fragment. Plasmid DNA from positive colonies was isolated and checked by Southern analysis. The insert of 1.25 kb in the smallest plasmid containing part of this *oppA* homologue (p15D4) was sequenced, which revealed the presence of an incomplete open reading frame (ORF3) that was 81% identical to *oppA* at the nucleotide level and, therefore, designated *appA*

(Fig. 1). The 5'-end (56 codons), an extra fragment towards the 3'-end (80 codons) and the region upstream of *appA* were obtained by inverse PCR (IPCR) on a 2.5-kb *Hind*III chromosomal fragment (Fig. 1). The region (1.8 kb) upstream of *appA* comprises a complete (ORF2) and an incomplete (ORF1) ORF. ORF1 comprises 228 codons and was designed *appB* although 49 codons are missing when DNA is aligned with *oppB*. The deduced protein AppB and OppB are 90.7% identical. ORF2 comprises 294 codons and the deduced protein, AppC, is 91.5% identical to OppC. *appC* is preceded by a putative ribosome-binding site (RBS) with a ΔG° of -12.8 kcal/mol. The intergenic region of *appC* and *appA* is 115 bp long and it contains consensus promoter sequences and a putative RBS with ΔG° of -9.4 kcal/mol (Van der Vossen et al., 1987). The sequence of the 3'-end of *appA* and the downstream region were obtained by IPCR on a 1.7-kb *Ssp*I chromosomal fragment (Fig. 1). The complete sequence of *appA* comprises 596 codons, which is four codons less than *oppA*, and the corresponding protein sequences are 82.8% identical. The region with the largest dissimilarity is position 235–276 (Fig. 2). Unlike for *oppA*, which is followed by the endopeptidase gene *pepO*, the region downstream of *appA* contains two inverted repeats (ΔG° of -10.5 and -5.7 kcal, respectively) that could function as a rho-independent terminator.

3.2. Inactivation of *appA* and functional characterization of disruption mutants

The gene *appA* was inactivated by plasmid integration via homologous recombination. One of the *appA*




Fig. 1. Schematic representation of the *opp* operon (Tynkkynen et al., 1993) and the sequenced chromosomal DNA region of *L. lactis* MG1363 containing the *opp* homologues, *appA*, *appC* and most of *appB*. The positions of putative promoter regions (P) and rho-independent terminators (T) are indicated. The restriction sites used for inverse PCR amplifications are indicated. The nucleotide sequence data will appear in the GenBank under accession number AF245305.

AppA	MKKLNITLLATTAVLATALLSACGSNQSSSTSTTKLKAGS	40
OppA	MnKlKvTLLAssvVLAatLLSACGSNQSSSTSTkKLKAGn	40
AppA	FDVAYKNPDKAIKGGDLKVAYQSDSPMKAEWWSGLENDAT	80
OppA	FDVAYqNPDKAIKGGnLKVAYQSDSPMKAqWLSGLsNDAT	80
AppA	FAAMASPAGGLDGIFFTNSAFKFINGGPANISLDDSAKTA	120
OppA	FAtMsgPgGGqDGlFFTdSgFKFIkGGaAdvaLDkesKTA	120
AppA	TITLRKDLKWSGDSQVTAKDYEFTYETIANPAYGSDRWTD	160
OppA	TITLRKDLKWSDGSeVTAKDYEFTYETIANPAYGSDRWTD	160
AppA	SLANIVGLSDYHTGKAKTISGITFPDGENGVKIKIQFKEM	200
OppA	SLANIVGLSDYHTGKAKTISGITFPDGENGVKIKvQFKEM	200
AppA	TPGMTQSGNGYFLETVPYQYLKDVAPKDLASSPKRQ.QT	239
OppA	kPGMTQSGNGYFLETVaPYQYLKDVAPKDLASSPKtttkp	240
AppA	ISHRSFQTANVVAGD.QSICTNPYYWA.KTKLT.LSPMNS	276
OppA	lvtgpFkpeNVVAGesikyvpNPYYWgeKpKlnsityevv	280
AppA	STNKSVAALSSGKYDFINSMVASQYKQVKNLKGKVLGQQ	316
OppA	STaKSVAALSSsKYDiINGMVSSQYKQVKNLKGKVLGQQ	320
AppA	ALYISLMYYNLGHYDTKNSINVQDRKTPLQDQSVRQAVAY	356
OppA	AmYISLMYYNLGHYDaKNSINVQDRKTPLQDQnVRQaigY	360
AppA	ARNVAEVDNKFNSGLATPANGLIPIPKQFTSSSVKGYEK	396
OppA	ARNVAEVDNKFNSGLsTPANsLIPPIFKQFTSSSVKGYEK	400
AppA	QNLDKANKLLDADGWKLNKSTGYREKDGKELSLVYAARQG	436
OppA	QdLDKANKLLDeDGWKLNKSTGYREKDGKELSLVYAARvG	440
AppA	DANQETIAQNYIQWKKIGVKVSLYNGKLMEFNSWDHMT	476
OppA	DANaETIAQNYIQWKKIGVKVSLYNGKLMEFNSWDHMT	480
AppA	TPPGADDWDITDGAWSLSSEPSQQDLFSAAAPYNIGHFND	516
OppA	TPPGAnDWDITDGsWSLaSEPSQQDLFSAAAPYNfGHFND	520
AppA	PQITNDLNDIDSTKSESATYRKAAFVKYQNDMNKKAYVVP	556
OppA	seITkDLNDIDSaKSEnPYRKAAFVKYQeDMNKKAYViP	560
AppA	T.FHLN.PPVNKRVLGMLTDYGDMLWSEIGVSSNKMATK	594
OppA	TnFmLNytPVNKRvVGMTLDYGaMntWSEIGVSSaKlATK	600

Fig. 2. Comparison of OppA and AppA proteins of *L. lactis* MG1363. The sequences were aligned with the DNAMAN program (Version 4.03). Identical residues in both sequences are in capital letters.

disruption mutants, designated *L. lactis* YS32, was selected and its phenotype was analyzed. It has been shown that the Opp system is responsible for the uptake of oligopeptides (>3 residues), and that the binding protein, OppA, is an essential component of this function (Kunji et al., 1996a). On the other hand, the existence of residual di- and tripeptide uptake in triple mutants, lacking functional Opp, DtpT and Dpp, suggests the presence of a fourth distinct uptake system (Sanz et al., 2001). Following these observations, the

appA disruption mutant was tested for its ability to utilize di- and tripeptides. Growth experiments were carried out in CDM containing all amino acids except an essential one that was supplied in the form of a di- or tripeptide (Poolman and Konings, 1988). Growth rates on several peptides of *L. lactis* YS32 and AG500, carrying the empty vector pAMP0 (Picón et al., 2000) were not significantly different (Fig. 3). Thus, *appA* does not seem to have a role in di- and tripeptide utilization.




Fig. 3. Growth of *L. lactis* strains in CDM containing a di- or tripeptide as the sole source of the essential amino acids. (A) CDM-Leu + 0.09 mM Leu-Leu: (●) AG500 (pAMP0) and (▲) YS32 (*appA*::p15D14dis); CDM-Leu + 0.18 mM Leu-Gly-Gly: (○) AG500 (pAMP0) and (△) YS32 (*appA*::p15D14dis); (B) CDM-Val + 0.28 mM Val-Val: (●) AG500 (pAMP0) and (▲) YS32 (*appA*::p15D14dis); CDM-Val + 0.58 mM Val-Gly-Gly: (○) AG500 (pAMP0) and (△) YS32 (*appA*::p15D14dis).

3.3. Expression of *appA* from a plasmid and role of *appA* in oligopeptide utilization

To determine whether or not *appA* encodes a functional binding protein, the gene was placed under the control of the constitutive lactococcal promoter *P32*. The expression of *oppA* and *appA* from the corresponding plasmids in *L. lactis* AMP2 ($\Delta oppA$) was quantified by Western analysis. Both proteins were expressed to similar levels (data not shown). Since the utilization of the pentapeptide Leu-enkephalin is characteristic for Opp activity, *L. lactis* AMP2 (pYSAprime) was tested for its ability to grow in CDM with Leu-enkephalin as the sole source of leucine. Growth rates of wild-type (MG1363) and the *oppA* deletion mutant AMP2, carrying either pAMP31 or pYSAprime, on different concentrations of Leu-enkephalin (0.18, 0.09 and 0.03 mM) were determined. As anticipated, the deletion mutant, *L. lactis* AMP2, was unable to grow

on any of these concentrations of Leu-enkephalin as the only source of leucine and each strain grew with similar rates when leucine was present (data not shown). The *appA* in pYSAprime complemented *oppA* with respect to Leu-enkephalin utilization, but the growth rate was much lower (Fig. 4). Growth rates of the wild-type (MG1363, data not shown) and the strain AMP2 (pAMP31) on Leu-enkephalin were similar ($\mu \sim 0.60 \text{ h}^{-1}$ at 0.18 mM Leu-enkephalin and $\sim 0.53 \text{ h}^{-1}$ at 0.09 mM Leu-enkephalin). Growth rates of AMP2 (pYSAprime) were considerably lower (0.21 h^{-1} at 0.18 mM Leu-enkephalin and 0.14 h^{-1} at 0.09 mM Leu-enkephalin), and growth was almost impaired at lower concentrations (0.03 mM) while MG1363 and AMP2 (pAMP31)




Fig. 4. Growth of *L. lactis* strains in CDM containing Leu-enkephalin as the sole source of the essential amino acid leucine. (A) 0.18 mM Leu-enkephalin; (B) 0.09 mM Leu-enkephalin; (C) 0.03 mM Leu-enkephalin. *L. lactis* strains: (●) AMP2; (■), AMP2 (pAMP31); (▲), AMP2 (pYSAprime).

grew to some extent. As these effects cannot be attributed to differences in expression of OppA and AppA, they most likely reflect the inefficient donation of the peptide by AppA to the translocator complex.

The absence of growth of *oppA* deletion mutants on Leu-enkephalin, and the complementation of *oppA* by *appA* when provided in trans suggest that the chromosomal copy of *appA* is not functional or poorly expressed. Western analysis of cell lysates of *L. lactis* AG500 confirms this notion using antibodies raised against OppA which also reacts to AppA (data not shown). Nevertheless, important structure–function relationships might be obtained from the dissimilarities between both proteins. The three dimensional structure of the peptide binding protein OppA of *Salmonella typhimurium* in complex with different ligands has made it possible (Sleigh et al., 1997; Tame et al., 1994, 1995). A first class of OppA specificity mutants has been constructed in *L. lactis* on the basis of multiple alignments with OppA from *S. typhimurium* (Picón et al., 2000). Future progress in defining residues of OppA critical for ligand binding can be made on the basis of sequence comparisons with the newly identified AppA.

Acknowledgements

This work was supported by the European Union (BIO-4-CT-960016) and ABON. Y. Sanz was a postdoctoral fellow of the Ministerio de Educación y Cultura (Spain). Michiel Hellendoorn is supported by the Ministry of Economic Affairs, the Ministry of Education, Culture, and Science, the Ministry of Agriculture, Nature Management and Fishery in the framework of an industrially relevant research program of the Netherlands Association of Biotechnology Centers in the Netherlands. Technical assistance of Mr. Rob Penninkhof is acknowledged.

References

- Alloung, G., de Philip, P., Claverys, J.-P., 1994. Three highly homologous membrane-bound lipoproteins participate in oligopeptide transport by the Ami system of the Gram-positive *Streptococcus pneumoniae*. *J. Mol. Biol.* 241, 44–58.
- Bimboim, H.C., Doly, J., 1979. A rapid alkaline extraction procedure for screening recombinant plasmid DNA. *Nucleic Acids Res.* 7, 1513–1523.
- Buist, G., Kok, J., Leenhouts, K.J., Dabrowska, M., Venema, G., Haandrikman, A.J., 1995. Molecular cloning and nucleotide sequence of the gene encoding the major peptidoglycan hydrolase of *Lactococcus lactis*, a muramidase needed for cell separation. *J. Bacteriol.* 177, 1554–1563.
- Ehrmann, M., Ehrle, R., Hofmann, E., Boos, W., Schlösser, A., 1998. The ABC maltose transporter. *Mol. Microbiol.* 29, 685–694.
- Hagting, A., Kunji, E.R.S., Leenhouts, K.J., Poolman, B., Konings, W.N., 1994. The di- and tripeptide transport protein of *Lactococcus lactis*: a new type of bacterial peptide transporter. *J. Biol. Chem.* 269, 11391–11399.
- Holo, H., Nes, Y.F., 1989. High-frequency transformation, by electroporation, of *Lactococcus lactis* subsp. *cremoris* grown with glycine in osmotically stabilized media. *Appl. Environ. Microbiol.* 53, 3119–3123.
- Kunji, E.R.S., Smid, E.J., Plapp, R., Poolman, B., Konings, W.N., 1993. Di-tripeptides and oligopeptides are taken up via distinct transport mechanism in *Lactococcus lactis*. *J. Bacteriol.* 175, 2052–2059.
- Kunji, E.R.S., Mierau, I., Hagting, A., Poolman, B., Konings, W.N., 1996a. The proteolytic systems of lactic acid bacteria. *Antonie van Leeuwenhoek* 70, 187–221.
- Kunji, E.R.S., Hagting, A., de Vries, C.J., Juillard, V., Haandrikman, A.J., Poolman, B., Konings, W.N., 1996b. Transport of β -casein-derived peptides by the oligopeptide transport system is a crucial step in the proteolytic pathway of *Lactococcus lactis*. *J. Biol. Chem.* 270, 1569–1574.
- Leenhouts, K.J., Venema, G., 1993. *Lactococcal plasmid vectors*. In: Hardy, K.G. (Ed.), *Plasmids: A Practical Approach*. IRL Press, Oxford, pp. 65–94.
- Leenhouts, K.J., Kok, J., Venema, G., 1989. Campbell-like integration of heterologous plasmid DNA into the chromosome of *Lactococcus lactis* subsp. *lactis*. *Appl. Environ. Microbiol.* 55, 394–400.
- Leenhouts, K.J., Kok, J., Venema, G., 1990. Stability of integrated plasmids in the chromosome of *Lactococcus lactis*. *Appl. Environ. Microbiol.* 56, 2726–2735.
- Marugg, J.D., Meijer, W., van Kranenburg, R., Laverman, P., Bruinenberg, P.G., de Vos, W.M., 1995. Medium-dependent regulation of proteinase gene expression in *Lactococcus lactis*: control of transcription initiation by specific dipeptides. *J. Bacteriol.* 177, 2982–2989.
- Meijer, W., Marugg, J.D., Hugenholtz, J., 1996. Regulation of proteolytic activity in *Lactococcus lactis*. *Appl. Environ. Microbiol.* 62, 156–161.
- Pearson, W., Lipman, D.J., 1988. Improved tools for biological sequence comparison. *Proc. Natl. Acad. Sci. U. S. A.* 85, 2444–2448.
- Picón, A., Kunji, E.R., Lanfermeijer, F.C., Konings, W.N., Poolman, B., 2000. Specificity mutants of the binding protein of the oligopeptide transport system of *Lactococcus lactis*. *J. Bacteriol.* 182, 1600–1608.
- Poolman, B., Konings, W.N., 1988. Relation of growth of *Strepto-*

- coccus lactis* and *Streptococcus cremoris* to amino acid transport. J. Bacteriol. 170, 700–707.
- Sambrook, J.E., Fritsch, F., Maniatis, T., 1989. Molecular Cloning: A Laboratory Manual, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Sanger, F., Nicklen, S., Coulson, A.R., 1977. DNA sequencing with chain-terminating inhibitors. Proc. Natl. Acad. Sci. U. S. A. 74, 5463–5467.
- Sanz, Y., Lanfermeijer, F.C., Konings, W.N., Poolman, B., 2000. Kinetics and structural requirements for the binding protein of the di-tripeptide transport system of *Lactococcus lactis*. Biochemistry 39, 4855–4862.
- Sanz, Y., Lanfermeijer, F.C., Renault, P., Bolotin, A., Konings, W.N., Poolman, B., 2001. Genetic and functional characterization of *dpp* genes encoding a dipeptide transport system in *Lactococcus lactis*. Arch. Microbiol. 175, 334–343.
- Sleigh, S.H., Tame, J.R.H., Dodson, E.J., Wilkinson, A.J., 1997. Peptide binding in OppA, the crystal structures of the periplasmic oligopeptide binding protein in the unliganded form and in complex with lysyllysine. Biochemistry 36, 9747–9758.
- Tame, J.R., Murshudov, G.N., Dodson, E.J., Neil, T.K., Dodson, G.G., Higgins, C.F., Wilkinson, A.J., 1994. The structural basis of the sequence-independent peptide binding by OppA protein. Science 264, 1578–1581.
- Tame, J.R., Dodson, E.J., Murshudov, G., Higgins, C.F., Wilkinson, A.J., 1995. The crystal structures of the oligopeptide-binding protein OppA complexed with tripeptide and tetrapeptide ligands. Structure 3, 1395–1406.
- Tynkkynen, S., Buist, G., Kunji, E., Kok, J., Poolman, B., Venema, G., Haandrikman, A., 1993. Genetic and biochemical characterization of the oligopeptide transport system of *Lactococcus lactis*. J. Bacteriol. 175, 7523–7532.
- Van der Vossen, J.M.B.M., Van der Lelie, D., Venema, G., 1987. Isolation and characterization of *Streptococcus cremoris* Wg2-specific promoters. Appl. Environ. Microbiol. 53, 2452–2457.
- Zabarovsky, E.R., Winberg, G., 1990. High efficiency of electroporation of ligated DNA into bacteria. Nucleic Acids Res. 18, 5912.